

12

EUROPEAN PATENT APPLICATION

21 Application number: **88120718.7**

51 Int. Cl.⁵: **C12P 7/18**

22 Date of filing: **12.12.88**

43 Date of publication of application:
20.06.90 Bulletin 90/25

64 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

71 Applicant: **UNILEVER NV**
Burgemeester s'Jacobplein 1 P.O. Box 760
NL-3000 DK Rotterdam(NL)

Applicant: **UNILEVER PLC**
Unilever House Blackfriars P.O. Box 68
London EC4P 4BQ(GB)

72 Inventor: **Gottschalk, G., Prof. Dr.**
Johann-Wolf-Strasse 35a
D-3412 Nörtenhardenberg(DE)
Inventor: **Averhoff, Beate, Dipl.-Biol.**
Osterlandweg 13
D-3400 Göttingen(DE)

74 Representative: **Hagemann, Heinrich, Dr.**
Dipl.-Chem. et al
Patentanwälte HAGEMANN & KEHL
Ismaninger Strasse 108 Postfach 860329
D-8000 München 86(DE)

54 Process for the microbiological preparation of 1,3-propane-diol from glycerol.

57 A process of the microbiological preparation of 1,3 propane diol from glycerol in growth media of suitable bacterial strains is described, accompanied by the addition of a cosubstrate in the form of a H-donor and the separation of the propane diol formed. It is characterized in that a) biomass is formed in a growth phase from the selected bacterial strain and accompanied by feeding with glycerol and, if necessary, whilst substantially excluding the H-donor until a stationary growth phase occurs and b) further glycerol and H-donor matched to the biomass are added to the resulting stationary cell suspension for increased 1,3-propane diol formation. This process makes it possible to produce 1,3-propane diol in a high yield from glycerol with a small amount of unobjectionable by-products in a batchwise manner or in continuous form, following immobilization.

EP 0 373 230 A1

THIS PAGE BLANK (USPTO)

THIS PAGE BLANK (USPTO)

Process for the microbiological preparation of 1,3-propane-diol from glycerol

The invention relates to a process for the microbiological preparation of 1,3-propane diol from glycerol in growth media of suitable bacterial strains, accompanied by the addition of a cosubstrate as the H-donor and the separation of the propane diol formed.

Processes for the preparation of 1,3-propane diol or trimethylene glycol (TMG) from glycerol are known. When performing these, essentially the following points must be taken into consideration. Bacterial strains able to convert glycerol into 1,3-propane diol are found e.g. in the species *Klebsiella* (cf. J. Bacteriol., 1982, 149, 413-419), *Citrobacter* (cf. J. Bacteriol., 1944, 39, 409-415), *Clostridium* (cf. Appl. Environ. Microbiol., 1987, 53, 639-643), *Ilyobacter* (cf. Arch. Microbiol., 1984, 140, 139-146) and *Lactobacillus* (cf. System. Appl. Microbiol., 1984, 5, 169-178). It cannot be excluded that further obligatory or optional ferments also form 1,3-propane diol as a fermentation product. Within the *Citrobacter* species, *C. freundii* is particularly well known as a 1,3-propane diol producer. Several strains (e.g. DSM 30039, 30040 and 30041) are available.

As a function of the bacterial strain used, it is possible to employ mineral media with glycerol or complex media with glycerol for the growth and associated production of 1,3-propane diol. The ambient conditions, such as the pH and temperature, are also dependent on the strain used. Preference is always given to anaerobic growth conditions, because the enzymes for the transformation of glycerol into 1,3-propane diol are only formed under anaerobic conditions. An adequate cobalt iron supply to the bacteria must be ensured, because the participating enzyme glycerol dehydratase contains coenzyme B₁₂ (cf. Arch. Biochem. Biophys., 1962, 97, 538-543).

Using the aforementioned bacterial strains, 1,3-propane diol can be produced batchwise or continuously. As a function of the species, different 1,3-propane diol yields are obtained in the batchwise anaerobic fermentation of glycerol: 78% for *Lactobacillus buchneri*, 61% für *Clostridium butylicum*, 40% for *Ilyobacter polytropus* and 50% for *Citrobacter freundii*. In connection with the *Lactobacillus* species, it must be borne in mind that a mixture of glycerol and glucose (2:1) and not glycerol alone can be fermented. As a function of the bacterial species during the fermentation of glycerol different by-product spectra appear, mainly acetate, butyrate, lactate, formate, succinate, ethanol, butanol and hydroxypropionate (in the case of *Ilyobacter polytropus*).

Whereas the aforementioned fermentations re-

late to cultures operating in a batchwise manner, in 1987 results were published for the first time concerning a continuous fermentation of glycerol for the production of 1,3-propane diol (cf. Appl. Environ. Microbiol., 1987, vol. 53, 639-643). Fermentation was carried out with *Clostridium butylicum* B 593, an obligatory ferment. 3.8% glycerol were added to the chemically defined medium and 97% thereof were reacted by the bacteria. It was also necessary to add 0.2% yeast extract to the medium. In the gas phase there was oxygen-free carbon dioxide, the residence time of the bacteria in the reactor was 10 hours and the temperature 35° C. Continuous fermentation gave the following product spectrum analyzed with HPLC and GC: 61% 1,3-propane diol, 6.6% acetate, 1.6% butyrate, 0.9% lactate, 0.6% butanol and 0.6% ethanol.

On the basis of the above statements, a 1,3-propane diol synthesis without by-products does not appear to be possible with bacteria, because part of the glycerol must be oxidized in order to obtain energy, whilst the reduction of glycerol to 1,3-propane diol is used for eliminating hydrogen which occurs. On lowering the pH-value from 6.5 to 4.9 the growth of bacteria and 1,3-propane diol production are reduced. An intermediate during the synthesis of 1,3-propane diol is 3-hydroxypropionaldehyde, whose production was also the aim of certain fermentation tests (cf. Appl. Environ. Microbiol., 1983, vol. 48, No. 1, 62-67). In the case of anaerobic fermentation of glycerol with *Klebsiella pneumoniae*, the speed of enzymatic reaction was displaced through the addition of semicarbazide hydrochloride. 3-hydroxypropionaldehyde production took place much more rapidly than the subsequent reduction to 1,3-propane diol and consequently led to a 3-hydroxypropionaldehyde accumulation. In the case of 30 g/l glycerol solutions, a 3-hydroxypropionaldehyde yield of 13.1 g/l was obtained.

The problem of the invention is to so improve the aforementioned process that it permits a simple, economic, rapid and continuous preparation of 1,3-propane diol from glycerol with a high yield and whilst substantially excluding environmentally prejudicial by-products.

According to the invention this problem is solved in that a) biomass is formed from the selected bacterial strain, accompanied by feeding with glycerol and, if necessary, with a substantial exclusion of the H-donor in a growth phase until a stationary growth phase is obtained and b) for increased 1,3-propane diol formation, to the corresponding stationary cell suspension are added

THIS PAGE BLANK (USPTO)

further glycerol and a H-donor matched to the biomass.

Thus, the essence of the inventive process is that initially in a process stage a) biomass is formed until a stationary growth phase or cell suspension has been obtained. It is necessary for most of the bacterial strains usable within the scope of the invention to substantially exclude a H-donor in said process stage a). This more particularly applies for the strains *Citrobacter freundii*, *Klebsiella pneumoniae*, *Clostridium autobutylicum* and *Clostridium butylicum*. However, several bacterial strains exist which, in the growth phase, grow in the presence of a H-donor and possibly even have a need for the latter, such as e.g. *Lactobacillus brevis* and *Lactobacillus buchneri*. However, the growth of most of the bacterial strains which can be used according to the invention is disturbed by the presence of a H-donor, so that in such cases the latter should be substantially excluded. This must also be borne in mind in connection with the qualitative evaluation of the glycerol-containing starting materials. Thus, if the glycerol starting material already contained a significant sugar fraction, such as glucose, then said fraction could be responsible for disturbing process stage a). Otherwise there are no significant limitations on the inventive process with regards to the glycerol starting material. Thus, it can e.g. be glycerol splitting water from high pressure splitting, or glycerol water from soap splitting, or glycerol water from the methanolysis of fats or oils, e.g. from fish oil, coconut oil or palm kernel oil.

The above explanations apply regarding the choice of the suitable bacterial strain. Representatives of the *Citrobacter* species, such as *Citrobacter freundii*, are used in a particularly advantageous manner. The strains DSM 30039, 30040 and 30041 are particularly suitable.

In process stage a), it is appropriate to adopt the following procedure: the description relating to *Citrobacter freundii* DSM 30040, but the remarks correspondingly apply to other bacterial strains. A mineral solution is prepared, in the manner known to the expert. However, as a special feature it contains a cobalt salt, such as in particular cobalt chloride in high concentration (250-550 µg CoCl₂/l) and an ammonium salt, such as in particular ammonium sulphate, as the growth-limiting N-source. As a function of the desired biomass concentration, the ammonium sulphate concentration can be approximately 0.4 to 1.5 g/l. Following the addition of glycerol (e.g. approximately 0.3 mol/l) and a preculture of strain DSM 30040, incubation takes place at approximately 37°C under anaerobic conditions. After about 12 to 20 hours (at 0.3 mol/l), the growing culture enters the stationary phase necessary for process stage b).

The concentration of the glycerol in process stage a) is not critical. It could be in a wide range, e.g. approximately 0.001 to 1 mol/l, although this range can also be exceeded. The range of approximately 0.1 to 0.4 mol/l is preferred, in particular approximately 0.3 mol/l.

It is not critical for performing process stage a) whether it is performed anaerobically or aerobically. It is preferably performed anaerobically, so that the enzymes particularly suitable for the desired conversion of glycerol into 1,3-propanediol are formed in an optimum manner. When process stage a) has allowed the desired biomass quantity to form, measures are taken to end the growth. This is preferably brought about in that the end of growth is controlled by adding a predetermined quantity of a phosphate or nitrogen source. Ammonium salts, such as ammonium sulphate, have proved to be particularly advantageous.

The inventive process can be performed both in a complex medium and in a mineral medium. Preference is given to the mineral medium, because it is less expensive, easier to handle and easier to dispose of at the end of the process. A mineral medium is understood to mean a medium, which only contains inorganic salts and additionally glycerol and possibly cysteine (as the reducing agent). The complex medium differs therefrom through a content of preferably approximately 1% yeast extract.

The biomass obtained according to process stage a) can be directly supplied to process stage b). It is also advantageous to concentrate the biomass by centrifuging or to immobilize it. Immobilization can take place in various ways, such as e.g. by calcium alginate, carageenan or polyurethanes. Immobilization preferably takes place with the readily available, commercial calcium alginate (Protanal-LF-20/60). The immobilized form of the biomass can e.g. be filled into a column, to which is continuously supplied the glycerol solution to be treated. This leads to the particular advantage of continuous process performance, the biomass being usable for weeks and without impairing its efficiency for obtaining 1,3-propanediol from glycerol. Tests have been carried out in which the biomass was active, without deterioration, for more than 42 days.

Glycerol is not essential for feeding the microorganisms for process stage b) and instead the combination glycerol/H-donor should be used. There are no significant limitations regarding the choice of the H-donor. The expert will instead choose the H-donor on the basis of his knowledge and the chosen bacterial strain. Generally, the H-donor can be constituted by mono-saccharides and disaccharides, particularly fructose and glucose. Glucose has proved particularly advantageous in

THIS PAGE BLANK (USPTO)

conjunction with *Citrobacter freundii* as the bacterial strain.

In order not to obtain premature inhibition in process stage b), a pH-value control is appropriate. The pH-value should be in the range approximately 5.5 to 8.5. The range approximately 6.5 to 7.5 and especially a value of approximately 7 are preferred. This gives optimum biotransformation conditions for the formation of 1,3-propane diol from glycerol.

There is a quantitative matching of the H-donor and glycerol for further optimization of the process. The precise numerical values for this are dependent on the nature of the H-donor and the nature of the chosen bacterial strain. No general guideline can be given. In the case of the preferably used bacterial strain *Citrobacter freundii* and an H-donor in the form of glucose, between the glucose and the glycerol the preferred molar ratio is approximately 1:7 to 1:8. The molar ratio between the glucose and the glycerol could generally be with respect to practical advantages approximately 1:1 to 1:25, in particular 1:5 to 1:10.

Process stage b) is also preferably performed in a mineral medium. In order to achieve the process objective in an optimum manner, a certain part is played by the glycerol concentration in the medium of process stage b). Excessive glycerol concentrations inhibit the participating enzyme systems. Inadequate glycerol concentrations lead to undesirably high processing costs. An approximately 0.2 to 1.5 molar glycerol solution and especially an approximately 1 molar glycerol solution is optimum for the medium treated during process stage b).

The invention can undergo further per se known modifications without any variation to the essence thereof. For example reference is made to the following:

Process stage a): modification of the mineral culture medium with regards to the concentrations of the individual components; concentration of the glycerol, whereby the source-dependent impurities can differ widely as regards nature and concentration; nature of the thorough mixing of the culture medium by stirring, pumping or gassing; provision of anaerobic conditions by reducing agents and inert gases.

Process stage b): concentration of the biomass; operating pH and temperature; nature or refeeding glycerol; nature and concentration of the H-donor.

The process according to the invention has numerous advantages compared with the hitherto known processes. As a result of the two-stage process performance, i.e. growth phase in a first process stage with transfer into a stationary phase and subsequent biotransformation phase, the 1,3-propane diol yield can be raised to over 90%. If the biomass formed in the first process stage is immo-

bilized and introduced into a column, this permits a long-term, continuous process performance, e.g. for more than 40 days. The glycerol is specifically converted into 1,3-propane diol and by-products are obtained in an unobjectionably small quantity. In certain cases they can be acetate and lactate, which can be simply removed and are not prejudicial to the environment. It is also possible to use an inexpensive nutrient medium in process stage a). In both stages, comparatively high glycerol concentrations can be chosen in the medium used. It is not necessary to abide by a particular purity of the starting glycerol. It is in fact also possible to use technical glycerols, provided that they contain no by-products disturbing the microorganisms (technical glycerols or raw glycerols). It may be the case that the technical glycerol in question is not suitable for process stage a), so that then purer glycerol starting materials would have to be used, whereas the less pure product can be used in the second process stage. This also represents an important advantage, because for the formation of the stationary phase or the biomass needed for the second process stage, a comparatively small glycerol quantity is required compared with the second process stage. This permits a more flexible performance of the process.

The invention is explained in greater detail hereinafter relative to an example.

Example

The course of process sequence a) is shown in fig. 1. *Citrobacter freundii* is allowed to grow in the following mineral medium under ammonium limitation: KH_2PO_4 , 6g/l; $\text{K}_2\text{HPO}_4 \cdot 3\text{H}_2\text{O}$, 14 g/l; $(\text{NH}_4)_2\text{SO}_4$, 0.04 g/l; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/l; CaCl_2 , 0.1 g/l; Cystein $\cdot \text{HCl}$, 0.1 g/l; trace element solution (1 l contains 5 g EDTA (disodium salt), 0.3 g $\text{H}_3\text{BO}_3 \cdot 4\text{H}_2\text{O}$, 0.03 g $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, 0.02 g $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 2 g $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.1 g $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 0.03 g $\text{NaMoO}_4 \cdot 2\text{H}_2\text{O}$, 0.02 g $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$, and 0.01 g $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$) (cf. Arch. Microbiol., 1966, 55, 245-256), 1 ml/l; CoCl_2 , 550 $\mu\text{g/l}$, glycerol 880 mmol/l. Following inoculation (20 ml preculture/1.2 l of main culture) incubation took place at 37° and a pH-value kept constant at 7.5. After 20 hours the culture passed into the stationary phase, in which glycerol is further reacted to 1,3-propane diol.

The reaction of glycerol to 1,3-propane diol in the presence of glucose as the H-donor through the cell suspension of *C. freundii* according to process stage b) is shown in fig. 2. In this case cells were suspended in potassium phosphate buffer of pH 7.5 in a concentration of 3 mg dry weight/ml. Incubation took place in the absence of oxygen (N_2 atmosphere) at 37° C after adding 240

THIS PAGE BLANK (USPTO)

mmol/l of glycerol and 42 mmol/l of glucos . The pH-value was held at 7.5 by means of titration with 2 N KOH. 191 mmol of 1,3-propane diol were formed from 211 mmol of glycerol, i.e. 91% on molar base and 75% on a weight base (maximum possible value 83%).

Explanations concerning figs. 1 and 4:

Fig. 1. The course of TMG production from glycerol in ammonium-limited, static culture in the presence of an 11 times increased Co^{2+} concentration. The TMG production and growth were followed in a 1.2 l culture in mineral medium with 0.04% ammonium sulphate, 880 mM glycerol and 550 $\mu\text{g Co}^{2+}/\text{l}$, the pH being kept constant at 7.5. Apart from TMG (∇), acetate (\square), ethanol (Δ), lactate (\bullet) and pyruvate (∇) were formed. The optical density is (\blacksquare) and glycerol (Δ).

Fig. 2 shows the course of TMG production through cell suspensions whilst maintaining the pH-value constant by means of KOH titration. TMG production from glycerol in the presence of 42 mM of glucose was followed through dilute cell suspensions (3 mg/ml) in KPO_4 buffer at pH 7.5. The pH-value was kept constant by means of titration with 2 N KOH. Glycerol (Δ) and TMG (\bullet) in the presence of KOH.

Fig. 3: Comparison of TMG production from glycerol through stationary cells at different constant pH-values. The tests were performed in 4 ml cell suspensions (protein concentration: 1.8 mg/ml) at constant pH-values of 6.6 (A), 7.0 (B), 7.9 (C) and 8.0 (D). The pH-value was kept constant by titrating with 0.6 N NaOH. Buffer used 248 mM KPO_4 (A-C) and 100 mM tris/HCl (D). Glycerol (Δ), TMG (\bullet), acetate (\blacksquare), ethanol (Δ).

Fig. 4: Comparison of TMG production from glycerol through stationary cells in the absence (A) or presence (B) of glucose. The course of TMG production from glycerol by cell suspension (protein concentration: 2.4 mg/ml) in KPO_4 buffer was followed at a constant pH-value of 7.5. Mixture B also contained 84 mM of glucose. Glycerol (Δ), TMG (\bullet), glucose (\square).

Claims

1. Process for the microbiological preparation of 1,3-propane diol from glycerol in growth media of suitable bacterial strains, accompanied by the addition of the cosubstrate in the form of a H-donor and separation of the propane diol formed, characterized in that a) biomass is formed from a growth phase from the selected bacterial strain and accompanied by feeding with glycerol and, if nec-

essary, whilst substantially excluding the H-donor until a stationary growth phase is obtained and b) further glycerol and a H-donor matched to the biomass are added to the stationary cell suspension formed, accompanied by increased 1,3-propane diol formation.

2. Process according to claim 1, characterized in that a sugar is added as a H-donor in process stage b).

3. Process according to claim 1 or 2, characterized in that a bacterial strain from the *Citrobacter* species is used.

4. Process according to claim 3, characterized in that as the bacterium of the *Citrobacter* species, use is made of a *Citrobacter freundii*.

5. Process according to at least one of the preceding claims, characterized in that anaerobic working takes place in process stage a).

6. Process according to at least one of the preceding claims, characterized in that anaerobic working takes place in process stage b).

7. Process according to at least one of the preceding claims, characterized in that a pH-value of approximately 6.5 to 8.5 is maintained in stages a) and b).

8. Process according to at least one of the preceding claims, characterized in that it is performed in a mineral medium.

9. Process according to at least one of the preceding claims, characterized in that the end of growth is controlled by the addition of a predetermined quantity of phosphate or nitrogen source.

10. Process according to claim 9, characterized in that an ammonium salt is used as the nitrogen source.



11. Process according to at least one of the preceding claims, characterized in that in process stage b) glycerol is initially present in an approximately 0.2 to 1.5 molar concentration.

12. Process according to at least one of the preceding claims, characterized in that in process stage a) the glycerol is initially present in an approximately 0.1 to 0.4 molar concentration.

13. Process according to at least one of the preceding claims 1 to 12, characterized in that the biomass obtained in process stage a) is immobilized and the medium of process stage b) is fed over the same.

14. Process according to claim 13, characterized in that immobilization is carried out with calcium alginate.

THIS PAGE BLANK (USPTO)



THIS PAGE BLANK (USPTO)




FIG. 3




FIG. 4

THIS PAGE BLANK (USPTO)



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 5)
D,A	CHEMICAL ABSTRACTS, vol. 106, no. 25, 22nd June 1987, page 537, abstract no. 212512a, Columbus, Ohio, US; C.W. FORSBERG: "Production of 1,3-propanediol from glycerol by Clostridium acetobutylicum and other Clostridium species", & APPL. ENVIRON. MICROBIOL. 1987, 53(4), 639-43 ---	1	C 12 P 7/18
D,A	CHEMICAL ABSTRACTS, vol. 101, 1984, page 340, abstract no. 166873r, Columbus, Ohio, US; H. SCHUETZ et al.: "Anaerobic reduction of glycerol to 1,3-propanediol by Lactobacillus brevis and Lactobacillus buchneri", & SYST. APPL. MICROBIOL. 1984, 5(2), 169-78 ---	1	
D,A	CHEMICAL ABSTRACTS, vol. 107, no. 7, 17th August 1987, page 333, abstract no. 54605x, Columbus, Ohio, US; E.A. JOHNSON et al.: "Klebsiella pneumoniae 1,3-propanediol:NAD oxidoreductase", & J. BACTERIOL. 1987, 169(5), 2050-4 ---	1	TECHNICAL FIELDS SEARCHED (Int. Cl. 5)
A	DE-A-3 336 051 (MAIZENA) * Claims * -----	1	C 12 P
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 10-08-1989	Examiner DELANGHE L. L. M.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

THIS PAGE BLANK (USPTO)



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 373 230 B1**

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: 17.02.93 (51) Int. Cl.⁵: **C12P 7/18**

(21) Application number: 88120718.7

(22) Date of filing: 12.12.88

(54) Process for the microbiological preparation of 1,3-propane-diol from glycerol.

(43) Date of publication of application:
20.06.90 Bulletin 90/25

(45) Publication of the grant of the patent:
17.02.93 Bulletin 93/07

(64) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(56) References cited:
DE-A- 3 336 051

CHEMICAL ABSTRACTS, vol. 106, no. 25,
22nd June 1987, page 537, abstract no.
212512a, Columbus, Ohio, US; C.W. FORS-
BERG: "Production of 1,3-propanediol from
glycerol by *Clostridium acetobutylicum* and
other *Clostridium* species", & **APPL. ENVI-
RON. MICROBIOL.** 1987, 53(4), 639-43

CHEMICAL ABSTRACTS, vol. 101, 1984, page
340, abstract no. 166873r, Columbus, Ohio,
US; H. SCHUETZ et al.: "Anaerobic reduction
of glycerol to 1,3-propanediol by *Lactobacil-
lus brevis* and *Lactobacillus buchneri*", &
SYST. APPL. MICROBIOL. 1984, 5(2), 169-78

CHEMICAL ABSTRACTS, vol. 107, no. 7, 17th
August 1987, page 333, abstract no. 54605x,

Columbus, Ohio, US; E.A. JOHNSON et al.:
"*Klebsiella pneumoniae* 1,3-propanediol:NAD
oxidoreductase", & **J. BACTERIOL.** 1987,
169(5), 2050-4

(73) Proprietor: **UNILEVER N.V.**
Weena 455
NL-3013 AL Rotterdam(NL)

Proprietor: **UNILEVER PLC**
Unilever House Blackfriars P.O. Box 68
London EC4P 4BQ(GB)

(72) Inventor: **Gottschalk, G., Prof. Dr.**
Johann-Wolf-Strasse 35a
W-3412 Nörtenhardenberg(DE)
Inventor: **Averhoff, Beate, Dipl.-Biol.**
Osterlandweg 13
W-3400 Göttingen(DE)

(74) Representative: **Hagemann, Heinrich, Dr.**
Dipl.-Chem. et al
**Patentanwälte HAGEMANN & KEHL Ismanin-
ger Strasse 108 Postfach 860329**
W-8000 München 86 (DE)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

EP 0 373 230 B1

Description

The invention relates to a process for the microbiological preparation of 1,3-propane diol from glycerol in growth media of suitable bacterial strains, accompanied by the addition of a cosubstrate as the H-donor and the separation of the propane diol formed.

Processes for the preparation of 1,3-propane diol or trimethylene glycol (TMG) from glycerol are known. When performing these, essentially the following points must be taken into consideration. Bacterial strains able to convert glycerol into 1,3-propane diol are found e.g. in the species *Klebsiella* (cf. J. Bacteriol., 1982, 149, 413-419), *Citrobacter* (cf. J. Bacteriol., 1944, 39, 409-415), *Clostridium* (cf. Appl. Environ. Microbiol., 1987, 53, 639-643), *Ilyobacter* (cf. Arch. Microbiol., 1984, 140, 139-146) and *Lactobacillus* (cf. System. Appl. Microbiol., 1984, 5, 169-178). It cannot be excluded that further obligatory or optional ferments also form 1,3-propane diol as a fermentation product. Within the *Citrobacter* species, *C. freundii* is particularly well known as a 1,3-propane diol producer. Several strains (e.g. DSM 30039, 30040 and 30041) are available.

As a function of the bacterial strain used, it is possible to employ mineral media with glycerol or complex media with glycerol for the growth and associated production of 1,3-propane diol. The ambient conditions, such as the pH and temperature, are also dependent on the strain used. Preference is always given to anaerobic growth conditions, because the enzymes for the transformation of glycerol into 1,3-propane diol are only formed under anaerobic conditions. An adequate cobalt iron supply to the bacteria must be ensured, because the participating enzyme glycerol dehydratase contains coenzyme B₁₂ (cf. Arch. Biochem. Biophys., 1962, 97, 538-543).

Using the aforementioned bacterial strains, 1,3-propane diol can be produced batchwise or continuously. As a function of the species, different 1,3-propane diol yields are obtained in the batchwise anaerobic fermentation of glycerol: 78% for *Lactobacillus buchneri*, 61% für *Clostridium butylicum*, 40% for *Ilyobacter polytropus* and 50% for *Citrobacter freundii*. In connection with the *Lactobacillus* species, it must be borne in mind that a mixture of glycerol and glucose (2:1) and not glycerol alone can be fermented. As a function of the bacterial species during the fermentation of glycerol different by-product spectra appear, mainly acetate, butyrate, lactate, formate, succinate, ethanol, butanol and hydroxypropionate (in the case of *Ilyobacter polytropus*).

Whereas the aforementioned fermentations relate to cultures operating in a batchwise manner, in

1987 results were published for the first time concerning a continuous fermentation of glycerol for the production of 1,3-propane diol (cf. Appl. Environ. Microbiol., 1987, vol. 53, 639-643). Fermentation was carried out with *Clostridium butylicum* B 593, an obligatory ferment. 3.8% glycerol were added to the chemically defined medium and 97% thereof were reacted by the bacteria. It was also necessary to add 0.2% yeast extract to the medium. In the gas phase there was oxygen-free carbon dioxide, the residence time of the bacteria in the reactor was 10 hours and the temperature 35°C. Continuous fermentation gave the following product spectrum analyzed with HPLC and GC: 61% 1,3-propane diol, 6.6% acetate, 1.6% butyrate, 0.9% lactate, 0.6% butanol and 0.6% ethanol.

On the basis of the above statements, a 1,3-propane diol synthesis without by-products does not appear to be possible with bacteria, because part of the glycerol must be oxidized in order to obtain energy, whilst the reduction of glycerol to 1,3-propane diol is used for eliminating hydrogen which occurs. On lowering the pH-value from 6.5 to 4.9 the growth of bacteria and 1,3-propane diol production are reduced. An intermediate during the synthesis of 1,3-propane diol is 3-hydroxypropionaldehyde, whose production was also the aim of certain fermentation tests (cf. Appl. Environ. Microbiol., 1983, vol. 46, No. 1, 62-67). In the case of anaerobic fermentation of glycerol with *Klebsiella pneumoniae*, the speed of enzymatic reaction was displaced through the addition of semicarbazide hydrochloride. 3-hydroxypropionaldehyde production took place much more rapidly than the subsequent reduction to 1,3-propane diol and consequently led to a 3-hydroxypropionaldehyde accumulation. In the case of 30 g/l glycerol solutions, a 3-hydroxypropionaldehyde yield of 13.1 g/l was obtained.

The problem of the invention is to so improve the aforementioned process that it permits a simple, economic, rapid and continuous preparation of 1,3-propane diol from glycerol with a high yield and whilst substantially excluding environmentally prejudicial by-products.

According to the invention this problem is solved in that a) biomass is formed from the selected bacterial strain, accompanied by feeding with glycerol and, if necessary, with a substantial exclusion of the H-donor in a growth phase until a stationary growth phase is obtained and b) for increased 1,3-propane diol formation, to the corresponding stationary cell suspension are added further glycerol and a H-donor matched to the biomass.

Thus, the essence of the inventive process is that initially in a process stage a) biomass is

formed until a stationary growth phase or cell suspension has been obtained. It is necessary for most of the bacterial strains usable within the scope of the invention to substantially exclude a H-donor in said process stage a). This more particularly applies for the strains *Citrobacter freundii*, *Klebsiella pneumoniae*, *Clostridium autobutylicum* and *Clostridium butylicum*. However, several bacterial strains exist which, in the growth phase, grow in the presence of a H-donor and possibly even have a need for the latter, such as e.g. *Lactobacillus brevis* and *Lactobacillus buchneri*. However, the growth of most of the bacterial strains which can be used according to the invention is disturbed by the presence of a H-donor, so that in such cases the latter should be substantially excluded. This must also be borne in mind in connection with the qualitative evaluation of the glycerol-containing starting materials. Thus, if the glycerol starting material already contained a significant sugar fraction, such as glucose, then said fraction could be responsible for disturbing process stage a). Otherwise there are no significant limitations on the inventive process with regards to the glycerol starting material. Thus, it can e.g. be glycerol splitting water from high pressure splitting, or glycerol water from soap splitting, or glycerol water from the methanolysis of fats or oils, e.g. from fish oil, coconut oil or palm kernel oil.

The above explanations apply regarding the choice of the suitable bacterial strain. Representatives of the *Citrobacter* species, such as *Citrobacter freundii*, are used in a particularly advantageous manner. The strains DSM 30039, 30040 and 30041 are particularly suitable.

In process stage a), it is appropriate to adopt the following procedure: the description relating to *Citrobacter freundii* DSM 30040, but the remarks correspondingly apply to other bacterial strains. A mineral solution is prepared, in the manner known to the expert. However, as a special feature it contains a cobalt salt, such as in particular cobalt chloride in high concentration (250-550 µg CoCl₂/l) and an ammonium salt, such as in particular ammonium sulphate, as the growth-limiting N-source. As a function of the desired biomass concentration, the ammonium sulphate concentration can be approximately 0.4 to 1.5 g/l. Following the addition of glycerol (e.g. approximately 0.3 mol/l) and a preculture of strain DSM 30040, incubation takes place at approximately 37°C under anaerobic conditions. After about 12 to 20 hours (at 0.3 mol/l), the growing culture enters the stationary phase necessary for process stage b).

The concentration of the glycerol in process stage a) is not critical. It could be in a wide range, e.g. approximately 0.001 to 1 mol/l, although this range can also be exceeded. The range of approxi-

mately 0.1 to 0.4 mol/l is preferred, in particular approximately 0.3 mol/l.

It is not critical for performing process stage a) whether it is performed anaerobically or aerobically. It is preferably performed anaerobically, so that the enzymes particularly suitable for the desired conversion of glycerol into 1,3-propane diol are formed in an optimum manner. When process stage a) has allowed the desired biomass quantity to form, measures are taken to end the growth. This is preferably brought about in that the end of growth is controlled by adding a predetermined quantity of a phosphate or nitrogen source. Ammonium salts, such as ammonium sulphate, have proved to be particularly advantageous.

The inventive process can be performed both in a complex medium and in a mineral medium. Preference is given to the mineral medium, because it is less expensive, easier to handle and easier to dispose of at the end of the process. A mineral medium is understood to mean a medium, which only contains inorganic salts and additionally glycerol and possibly cystein (as the reducing agent). The complex medium differs therefrom through a content of preferably approximately 1% yeast extract.

The biomass obtained according to process stage a) can be directly supplied to process stage b). It is also advantageous to concentrate the biomass by centrifuging or to immobilize it. Immobilization can take place in various ways, such as e.g. by calcium alginate, carageenan or polyurethanes. Immobilization preferably takes place with the readily available, commercial calcium alginate (Protanal-LF-20/60). The immobilized form of the biomass can e.g. be filled into a column, to which is continuously supplied the glycerol solution to be treated. This leads to the particular advantage of continuous process performance, the biomass being usable for weeks and without impairing its efficiency for obtaining 1,3-propane diol from glycerol. Tests have been carried out in which the biomass was active, without deterioration, for more than 42 days.

Glycerol is not essential for feeding the microorganisms for process stage b) and instead the combination glycerol/H-donor should be used. There are no significant limitations regarding the choice of the H-donor. The expert will instead choose the H-donor on the basis of his knowledge and the chosen bacterial strain. Generally, the H-donor can be constituted by mono-saccharides and disaccharides, particularly fructose and glucose. Glucose has proved particularly advantageous in conjunction with *Citrobacter freundii* as the bacterial strain.

In order not to obtain premature inhibition in process stage b), a pH-value control is appropriate.

The pH-value should be in the range approximately 5.5 to 8.5. The range approximately 6.5 to 7.5 and especially a value of approximately 7 are preferred. This gives optimum biotransformation conditions for the formation of 1,3-propane diol from glycerol.

There is a quantitative matching of the H-donor and glycerol for further optimization of the process. The precise numerical values for this are dependent on the nature of the H-donor and the nature of the chosen bacterial strain. No general guideline can be given. In the case of the preferably used bacterial strain *Citrobacter freundii* and an H-donor in the form of glucose, between the glucose and the glycerol the preferred molar ratio is approximately 1:7 to 1:8. The molar ratio between the glucose and the glycerol could generally be with respect to practical advantages approximately 1:1 to 1:25, in particular 1:5 to 1:10.

Process stage b) is also preferably performed in a mineral medium. In order to achieve the process objective in an optimum manner, a certain part is played by the glycerol concentration in the medium of process stage b). Excessive glycerol concentrations inhibit the participating enzyme systems. Inadequate glycerol concentrations lead to undesirably high processing costs. An approximately 0.2 to 1.5 molar glycerol solution and especially an approximately 1 molar glycerol solution is optimum for the medium treated during process stage b).

The invention can undergo further per se known modifications without any variation to the essence thereof. For example reference is made to the following:

Process stage a): modification of the mineral culture medium with regards to the concentrations of the individual components; concentration of the glycerol, whereby the source-dependent impurities can differ widely as regards nature and concentration; nature of the thorough mixing of the culture medium by stirring, pumping or gassing; provision of anaerobic conditions by reducing agents and inert gases.

Process stage b): concentration of the biomass; operating pH and temperature; nature or refeeding glycerol; nature and concentration of the H-donor.

The process according to the invention has numerous advantages compared with the hitherto known processes. As a result of the two-stage process performance, i.e. growth phase in a first process stage with transfer into a stationary phase and subsequent biotransformation phase, the 1,3-propane diol yield can be raised to over 90%. If the biomass formed in the first process stage is immobilized and introduced into a column, this permits a long-term, continuous process performance, e.g. for more than 40 days. The glycerol is specifically converted into 1,3-propane diol and by-products

are obtained in an unobjectionably small quantity. In certain cases they can be acetate and lactate, which can be simply removed and are not prejudicial to the environment. It is also possible to use an inexpensive nutrient medium in process stage a). In both stages, comparatively high glycerol concentrations can be chosen in the medium used. It is not necessary to abide by a particular purity of the starting glycerol. It is in fact also possible to use technical glycerols, provided that they contain no by-products disturbing the microorganisms (technical glycerols or raw glycerols). It may be the case that the technical glycerol in question is not suitable for process stage a), so that then purer glycerol starting materials would have to be used, whereas the less pure product can be used in the second process stage. This also represents an important advantage, because for the formation of the stationary phase or the biomass needed for the second process stage, a comparatively small glycerol quantity is required compared with the second process stage. This permits a more flexible performance of the process.

The invention is explained in greater detail hereinafter relative to an example.

Example

The course of process sequence a) is shown in fig. 1. *Citrobacter freundii* is allowed to grow in the following mineral medium under ammonium limitation: KH_2PO_4 , 6g/l; $\text{K}_2\text{HPO}_4 \cdot 3\text{H}_2\text{O}$, 14 g/l; $(\text{NH}_4)_2\text{SO}_4$, 0.04 g/l; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/l; CaCl_2 , 0.1 g/l; Cystein $\cdot \text{HCl}$, 0.1 g/l; trace element solution (1 l contains 5 g EDTA (disodium salt), 0.3 g $\text{H}_3\text{BO}_3 \cdot 4\text{H}_2\text{O}$, 0.03 g $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, 0.02 g $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 2 g $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.1 g $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 0.03 g $\text{NaMoO}_4 \cdot 2\text{H}_2\text{O}$, 0.02 g $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$, and 0.01 g $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$) (cf. Arch. Microbiol., 1966, 55, 245-256), 1 ml/l; CoCl_2 , 550 $\mu\text{g/l}$, glycerol 880 mmol/l. Following inoculation (20 ml preculture/1.2 l of main culture) incubation took place at 37° and a pH-value kept constant at 7.5. After 20 hours the culture passed into the stationary phase, in which glycerol is further reacted to 1,3-propane diol.

The reaction of glycerol to 1,3-propane diol in the presence of glucose as the H-donor through the cell suspension of *C. freundii* according to process stage b) is shown in fig. 2. In this case cells were suspended in potassium phosphate buffer of pH 7.5 in a concentration of 3 mg dry weight/ml. Incubation took place in the absence of oxygen (N_2 atmosphere) at 37° C after adding 240 mmol/l of glycerol and 42 mmol/l of glucose. The pH-value was held at 7.5 by means of titration with 2 N KOH. 191 mmol of 1,3-propane diol were formed from 211 mmol of glycerol, i.e. 91% on molar base and 75% on a weight base (maximum

possible value 83%).

Explanations concerning figs. 1 and 4:

Fig. 1. The course of TMG production from glycerol in ammonium-limited, static culture in the presence of an 11 times increased Co^{2+} concentration. The TMG production and growth were followed in a 1.2 l culture in mineral medium with 0.04% ammonium sulphate, 880 mM glycerol and 550 $\mu\text{g Co}^{2+}/\text{l}$, the pH being kept constant at 7.5. Apart from TMG (∇), acetate (\square), ethanol (Δ), lactate (\bullet) and pyruvate (∇) were formed. The optical density is (\blacksquare) and glycerol (Δ).

Fig. 2 shows the course of TMG production through cell suspensions whilst maintaining the pH-value constant by means of KOH titration. TMG production from glycerol in the presence of 42 mM of glucose was followed through dilute cell suspensions (3 mg/ml) in KPO_4 buffer at pH 7.5. The pH-value was kept constant by means of titration with 2 N KOH. Glycerol (Δ) and TMG (\bullet) in the presence of KOH.

Fig. 3: Comparison of TMG production from glycerol through stationary cells at different constant pH-values. The tests were performed in 4 ml cell suspensions (protein concentration: 1.8 mg/ml) at constant pH-values of 6.6 (A), 7.0 (B), 7.9 (C) and 8.0 (D). The pH-value was kept constant by titrating with 0.6 N NaOH. Buffer used 248 mM KPO_4 (A-C) and 100 mM tris/HCl (D).

Glycerol (Δ), TMG (\bullet), acetate (\blacksquare), ethanol (Δ).

Fig. 4: Comparison of TMG production from glycerol through stationary cells in the absence (A) or presence (B) of glucose. The course of TMG production from glycerol by cell suspension (protein concentration: 2.4 mg/ml) in KPO_4 buffer was followed at a constant pH-value of 7.5. Mixture B also contained 84 mM of glucose. Glycerol (Δ), TMG (\bullet), glucose (\square).

Claims

1. Process for the microbiological preparation of 1,3-propane diol from glycerol in growth media of suitable bacterial strains, accompanied by the addition of a cosubstrate in the form of an H-donor and separation of the 1,3-propanediol formed, characterized in that

a) a biomass is formed by culturing the selected bacterial strain in the growth phase in a growth medium containing glycerol, but with the substantial exclusion of any H-donor;

b) the bacterial cells are transferred to a stationary phase and biotransformation is

effect d by adding further glycerol and an H-donor matched to the biomass, accompanied by increased 1,3-propane diol formation, this being the main stage of its preparation.

2. Process according to claim 1, characterized in that a sugar is added as H-donor in step b).

3. Process according to claim 1 or 2, characterized in that the bacterial strain is a strain from the *Citrobacter* genus.

4. Process according to claim 3, characterized in that as the bacterial strain is a strain of *Citrobacter freundii*.

5. Process according to claim 1 or 2, characterized in that an H-donor is added to the growth phase if the selected bacterial strain grows in the presence of an H-donor or has a need for it.

6. Process according to any preceding claim, characterized in that anaerobic working takes place in step a).

7. Process according to any preceding claim, characterized in that anaerobic working takes place in step b).

8. Process according to any preceding claim, characterized in that a pH-value of approximately 6.5 to 8.5 is maintained in steps a) and b).

9. Process according to any preceding claim, characterized in that steps a) and b) are performed in a mineral medium.

10. Process according to any preceding claim, characterized in that the transfer of the growing cells to the stationary phase is performed by the addition of a predetermined quantity of phosphate or nitrogen source.

11. Process according to claim 10, characterized in that an ammonium salt is used as the nitrogen source or a potassium dihydrogen phosphate is used as the phosphate source.

12. Process according to any preceding claim, characterized in that glycerol is initially present in step b) in approximately 0.2 to 1.5 molar concentration.

13. Process according to any preceding claim, characterized in that glycerol is initially present

in step a) in approximately 0.1 to 0.4 molar concentration.

14. Process according to any preceding claim characterized in that said biomass obtained in step a) is immobilized before step b). 5
15. Process according to any preceding claim, characterized in that said immobilization is carried out with calcium alginate. 10

Patentansprüche

1. Verfahren zur mikrobiologischen Herstellung von 1,3-Propandiol aus Glycerin in Wachstumsmedien geeigneter Bakterienstämme unter Zusatz eines Cosubstrates in Form eines H-Donators und Abtrennung des gebildeten 1,3-Propandiols, dadurch gekennzeichnet, daß
 - a) eine Biomasse durch Kultivieren des gewählten Bakterienstammes in der Wachstumsphase in einem Glycerin enthaltenden Wachstumsmedium, jedoch unter weitgehendem Ausschluß von jeglichem H-Donator, gebildet wird,
 - b) die Bakterienzellen in eine stationäre Phase überführt werden und die Biotransformation bewirkt wird, indem man weiteres Glycerin sowie einen auf die Biomasse abgestimmten H-Donator zufügt, was von einer die Hauptstufe seiner Herstellung darstellenden verstärkten Bildung von 1,3-Propandiol begleitet ist.
 15
2. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß in Stufe b) ein Zucker als H-Donator zugefügt wird. 20
3. Verfahren nach Anspruch 1 oder 2, dadurch gekennzeichnet, daß der Bakterienstamm ein Stamm der Gattung Citrobacter ist. 25
4. Verfahren nach Anspruch 3, dadurch gekennzeichnet, daß der Bakterienstamm ein Stamm von Citrobacter freundii ist. 30
5. Verfahren nach Anspruch 1 oder 2, dadurch gekennzeichnet, daß ein H-Donator zu der Wachstumsphase gegeben wird, wenn der gewählte Bakterienstamm in Gegenwart eines H-Donators wächst oder dessen bedarf. 35
6. Verfahren nach irgendeinem der vorhergehenden Ansprüche, dadurch gekennzeichnet, daß die Stufe a) anaerob abläuft. 40
7. Verfahren nach irgendeinem der vorhergehenden Ansprüche, dadurch gekennzeichnet, daß 45

die Stufe b) anaerob abläuft.

8. Verfahren nach irgendeinem der vorhergehenden Ansprüche, dadurch gekennzeichnet, daß in den Stufen a) und b) ein pH-Wert von etwa 6,5 bis 8,5 aufrechterhalten wird.
9. Verfahren nach irgendeinem der vorhergehenden Ansprüche, dadurch gekennzeichnet, daß die Stufen a) und b) in einem mineralischen Medium durchgeführt werden.
10. Verfahren nach irgendeinem der vorhergehenden Ansprüche, dadurch gekennzeichnet, daß die Überführung der wachsenden Zellen in die stationäre Phase durch Zugabe einer vorherbestimmten Menge einer Phosphat- oder Stickstoffquelle erfolgt.
11. Verfahren nach Anspruch 10, dadurch gekennzeichnet, daß ein Ammoniumsalz als Stickstoffquelle oder ein Kaliumdihydrogenphosphat als Phosphatquelle verwendet wird.
12. Verfahren nach irgendeinem der vorhergehenden Ansprüche, dadurch gekennzeichnet, daß Glycerin anfangs in Stufe b) in einer molaren Konzentration von etwa 0,2 bis 1,5 vorliegt.
13. Verfahren nach irgendeinem der vorhergehenden Ansprüche, dadurch gekennzeichnet, daß Glycerin anfangs in Stufe a) in einer molaren Konzentration von etwa 0,1 bis 0,4 vorliegt.
14. Verfahren nach irgendeinem der vorhergehenden Ansprüche, dadurch gekennzeichnet, daß die in Stufe a) erhaltene Biomasse vor der Stufe b) immobilisiert wird.
15. Verfahren nach irgendeinem der vorhergehenden Ansprüche, dadurch gekennzeichnet, daß die Immobilisierung mit Calciumalginat erfolgt. 50

Revendications



1. Procédé de préparation microbiologique de 1,3-propanediol à partir du glycérol dans des milieux de croissance de souches bactériennes appropriées, avec addition d'un cosubstrat sous forme d'un donneur de H et séparation du 1,3-propanediol formé, caractérisé en ce que:
 - a) une biomasse est formée par culture de la souche bactérienne choisie dans la phase de croissance au sein d'un milieu de croissance contenant du glycérol, mais avec exclusion pratique du donneur de H ;
 - b) les cellules bactériennes sont transférées à une phase stationnaire et on effectue la 55

biotransformation par addition d'un supplément de glycérol et d'un donneur de H₂ en concordance avec la biomasse, ce qui s'accompagne d'une plus forte formation de 1,3-propanediol, ce stade constituant le stade principal de la préparation.

2. Procédé selon la revendication 1, caractérisé en ce qu'on ajoute un sucre en qualité de donneur de H₂ au stade b). 10
3. Procédé selon la revendication 1 ou 2, caractérisé en ce que la souche bactérienne est une souche provenant du genre *Citrobacter*. 15
4. Procédé selon la revendication 3, caractérisé en ce que la souche bactérienne est une souche de *Citrobacter freundii*. 20
5. Procédé selon la revendication 1 ou 2, caractérisé en ce qu'on ajoute le donneur de H₂ à la phase de croissance si la souche bactérienne choisie pousse en présence d'un donneur de H₂ ou nécessite celui-ci dans ce but. 25
6. Procédé selon l'une quelconque des revendications précédentes, caractérisé en ce qu'au stade a), le procédé est anaérobie. 30
7. Procédé selon l'une quelconque des revendications précédentes, caractérisé en ce qu'au stade b), le procédé est anaérobie. 35
8. Procédé selon l'une quelconque des revendications précédentes caractérisé en ce que dans les stades a) et b), on maintient le pH à environ 6,5 à 8,5. 40
9. Procédé selon l'une quelconque des revendications précédentes, caractérisé en ce qu'on exécute les stades a) et b) dans un milieu minéral. 45
10. Procédé selon l'une quelconque des revendications précédentes caractérisé en ce qu'on effectue le transfert des cellules en croissance à la phase stationnaire par addition d'une quantité prédéterminée d'une source de phosphate ou d'azote. 50
11. Procédé selon la revendication 10, caractérisé en ce qu'on utilise un sel d'ammonium comme source d'azote ou un dihydrogène-phosphate de potassium comme source de phosphate. 55
12. Procédé selon l'une quelconque des revendications précédentes, caractérisé en ce que le glycérol est initialement présent au stade b) en

une concentration molaire d'environ 0,2 à 1,5.

13. Procédé selon l'une quelconque des revendications précédentes, caractérisé en ce que le glycérol est initialement présent au stade a) en une concentration molaire d'environ 0,1 à 0,4.
14. Procédé selon l'une quelconque des revendications précédentes, caractérisé en ce qu'on immobilise la biomasse obtenue au stade a) avant le stade b).
15. Procédé selon l'une quelconque des revendications précédentes, caractérisé en ce qu'on effectue ladite immobilisation avec de l'alginate de calcium.






FIG. 3




FIG. 4

THIS PAGE BLANK (USPTO)